The Excel Math Competition



May 2025 (Calculus Exam)

- 1. This is a 60 minute individual exam.
- 2. No collaboration or external devices (like a calculator) are allowed.
- 3. The first 10 questions are worth [5] points each and are multiple choice.
- 4. The last 5 questions are worth [10] points each and are short response.
- 5. Each of the final 5 questions have answers which are positive integers between 000 and 999, inclusive.
- 6. The questions are arranged in roughly ascending difficulty.
- 7. If you believe a question is seriously flawed, or have an answer which is not one of the listed answers, there will be a 10-minute dispute period after the test, after which no disputes will be accepted.
- 8. In the event of a dispute, **leave the question blank** and let your proctor know after the testing time ends.
- 9. All disputes will be considered on an individual-by-individual basis, so no student will receive credit if they did not submit a dispute, except for in the case of a question being thrown out.

Test writer: David Zapata Joseph Levenston

Test Editor: David Zapata **Problem Contributors:** David Zapata Joseph Levenston Siyuan Du Sponsors:





$1 \quad [5]$

Welcome to the Excel Academe Math Competition! This is our first time doing a calculus test ever, so congratulations on being a part of this historical moment! We hope everyone did good on their AP exams! The following equates to a simplified fraction of the form $\frac{m}{k}$. What is m + k?

$2 \quad [5]$

Find the slope of the normal line of $x^3 + 2x + 5$ at x = 1

a)
$$-\frac{1}{5}$$
 b) $\frac{1}{5}$ c) -5 d) 5 e) 0

3 [5]

David is running on 30 minutes of sleep again and has Calculus homework due at 11:59PM (it is currently 11:50PM), solve this integral so David can get a 100% on his homework.

$4 \quad [5]$

If $f(x) = e^{3x-3} + 5x^2 - 6$, find the derivative of $f^{-1}(x)$ at x = 0a) $-\frac{1}{13}$ b) $\frac{1}{13}$ c) 13 d) -13 e) 0

5 [5]

$$\int \frac{\sqrt{x^2 + 16}}{x^4} = -\frac{(x^{\alpha} + \beta)^{\gamma}}{\lambda \cdot x^{\rho}} + C$$

What is $\alpha + \beta + 2 \cdot \gamma + \lambda + \rho$? a) 72 b) 81 c) 90 d) 99 e) 108 $6 \quad [5]$

a) 0 b)
$$\frac{1}{2007}$$
 c) $\frac{1}{\sqrt{n}} x^{5x^2 + 18x + 2008} dx = ?$
a) 0 b) $\frac{1}{2007}$ c) $\frac{1}{2008}$ d) $\frac{1}{2009}$ e) DNE

 $7 \quad [5]$

An important concept to understand in the study of classical physics is the center of mass of an object with an non-uniform density. For example, if we were given a function for r in terms of mass in the x direction, we can find the center of mass in the x direction by using the formula $x_{cm} = \frac{\int r \cdot dm}{\int dm}$, Suppose we had a strictly 1 dimensional linear rod that is 5 meters long and has a function that gives the density at a specified length which is $\lambda(x) = x^2 + 3\sqrt{x}$ where x = 0 is the to the left of the rod. Let x_{cm} be the center of mass of this rod and let M be the total mass of this rod. $|x_{cm}| - M = b\sqrt{5} + \frac{m}{n}$. What is b + m + n?

8 [5]

In statistics, a probability density function (PDF) allows us to calculate the probability of a continuous random variable falling within a specific interval by finding the area under the curve of the PDF between the chosen bounds and the x-axis. The probability density function of the standard normal distribution (a normal distribution with a mean of 0 and a standard deviation of 1) is:

$$\frac{1}{\sqrt{2\pi}} \cdot e^{-\frac{x^2}{2}}$$

Using this knowledge, answer the following:

$$\int_{-\infty}^{\infty} e^{-x^2} dx$$
a) $\sqrt{\frac{\pi}{2}}$ b) $\sqrt{\frac{3\pi}{2}}$ c) $\sqrt{2\pi}$ d) \sqrt{e} e) $\sqrt{\pi}$

9 [5]

$$\int_{0}^{1} e^{-x^{2}} dx + \int_{-\sqrt{e}}^{\sqrt{e}} \frac{\sin(x)}{1 + x^{2} + \cos(x)} dx + \int_{1}^{e^{-1}} \sqrt{\ln(\frac{1}{x})} dx = e^{k}$$

What is k?

a) -2 b) -1 c) 0 d) 1 e) 2

10 [5]

a)
$$\frac{2^{e+2}}{e+2}$$
 b) $\frac{2^{e+2}}{e}$ c) $\frac{2^e}{e}$ d) $\frac{2^e+1}{e}$ e) $\frac{2^e+1}{e-2}$

$11 \quad [10]$

If the area of the inner loop of the polar curve $r = 1 + 2\cos\theta$ is equal to $\pi - \frac{A\sqrt{B}}{C}$ where the fraction is in simplest form and properly rationalized, what is $A \cdot B \cdot C$?

$$12 \quad [10]$$

If
$$I_n = \int_{\frac{n}{2}}^{\infty} {n \choose x} dx$$
, then find $\sum_{n=0}^{\infty} \frac{1}{I_n}$

13 [10]

$$\int_{1}^{2} \frac{3x^{4} + 4x + 6x^{-1}}{x^{6} + 3x^{5} + x^{2} + 1} dx = \ln(\frac{A}{B})$$

What is A + B given that $\frac{A}{B}$ is a fraction in simplest form.

$14 \quad [10]$

$$S = \sum_{n=1}^{\infty} \frac{\cos(n)}{n}$$

What is $\lfloor 100S \rfloor$?

15 [10]

$$L = \int_{-1}^{1} \sqrt{\frac{x-1}{x+1}} \cdot \arccos(\mathbf{x})$$

If $L = \frac{\pi^a}{b} - c$, what is a^{b^c}